

Docket No.: 7049405002
Customer No. 23,639

Application Serial No. 09/575,377

REMARKS

BEST AVAILABLE COPY

The communication of March 28, 2006, refused entry of the amendment filed with the RCE (Jan. 23, 2006) on grounds that the claims would be directed to a distinct invention from a method from "identifying one or more ion channels of a cell that may be affected by a test substance. Accordingly, the claims submitted herewith are directed to that invention. The amendment filed August 22, 2005 was not entered, so the present changes are based on the claims as they appear in the Amendment filed November 26, 2004. Claims 51 - 72 remain active in the case. Claim 73 has been cancelled as redundant in view of amended claim 51.

Support for classifying a test substance in one or more functional categories based on reference to a data library formed from known compounds is found on page 30, lines 11 - 12; page 31, lines 7 - 15; page 32, Table 1 and page 33, lines 1 - 17.

Serum-free culture medium is supported on page 26, lines 4 - 6 ("The neurons are grown ... in serum-free media. The cells are cultured in defined media...").

An intervening layer acting as a high-impedance seal is supported in the specification at page 11, lines 6 - 12.

Measurement of changes in one or more characteristics selected from after potential, time to cessation of activity, frequency, amplitude, shape, spike rate, or time constant are described on page 40, lines 14 - 31, and this subject matter was also in claim 53.

Deconvoluting changes to identify one or more ion channels affected by the test substance is supported on page 7, lines 13 - 28 and page 10, lines 4 - 16. Please note we specifically define deconvolution as not including spectral analysis on page 13, lines 1 - 5. Therefore, the term "deconvolution" does not read on Fourier Transform analysis.

Our initial description of "functional categories" came from the literature (see ref Riley et al. in the application as well as page 6 lines 17-18) and it is a method of classifying related genes and their gene products and thus the pathways they enable. We have shown differences in Action Potential peak shape with the data included in the application, see Figure 5.

We have since demonstrated the application in our paper "Toxin Detection Based on Action Potential Peak Shape Analysis using a Realistic Mathematical Model of Differentiated NG106-15 Cells" that has been accepted and is in press in the journal Biosensors and Bioelectronics (Mohan, et al., submitted previously).

Docket No.: 7049405002
Customer No. 23,639

Application Serial No. 09/575,377

Table 1 on page 32 of the application also further illustrates our ability to group the effects of different compounds into different "functional categories."

As discussed on page 40 lines 21-32 and page 41 lines 1-4 of the application, deconvolution of an action potential implies that some prior knowledge of the factors contributing to the shapes are known. It is well known that an action potential requires the contributions of ion fluxes through at least three distinct ion channels. Varying the contributions of these ion fluxes from these channels has been shown to vary the shape of the action potential, and this is illustrated in Figures 5 and 6A of the application. Thus, to deconvolute an action potential into its components and relate them to pathways or functional categories inside a cell requires some prior knowledge of the biology of the system (i.e., a data base of biological test data on the subject cell culture) in order to perform the deconvolution step and identify cell pathways. The Mohan et al. paper illustrates some of our efforts in this area and an implementation of the algorithm shown in Figure 10 of the patent application. This algorithm has been used to deconvolute one of the examples given in Figure 5 of the application and is shown in Figure A below.




Figure A. Result of parameter fitting to action potential shapes before (left panel) and after (right panel) the application of 1 mM Ricin. Action potentials were recorded from differentiated NG108-15 cells using whole-cell patch clamp method. Action potentials were fitted (dotted lines) with a computer model of NG108-15 cells established based on the measurement of ion channel parameters.

Control Activation

Inactivation

Docket No.: 7049405002
Customer No. 23,639

Application Serial No. 09/575,377

		V_{rev}	z	$V_{1/2}$	g	A	z	$V_{1/2}$	g	A
Na	140.00	100.00	6.00	-59.93	-0.45	1.80	-7.48	-70.36	0.42	8.00
K	26.00	80.00	8.00	-10.00	-0.60	10.00				
Ca	11.00	32.00	4.10	-11.00	-0.67	0.47				
Ricin										
Na	75.00	100.00	6.00	-59.93	-0.42	3.00	-7.48	-70.36	0.42	12.00
K	26.00	80.00	8.00	-10.00	-0.60	15.00				
Ca	11.00	32.00	4.10	-11.00	-0.67	0.80				

Table 1. Effect of 1 mM Ricin on the action potential parameters. Action potential parameters were obtained by fitting the parameters of our NG108-15 cell model to the experimental data. 1 mM Ricin caused a significant slow-down of the sodium channel kinetics (parameter A) and a decrease of sodium conductances in the membrane (parameter g). The effect of Ricin on the other ion channels were not so profound, partly because potassium and calcium channels are activated at higher membrane potentials, so their contribution to the Ricin-modified action potential is less (the amplitude of the action potential is smaller, barely reaching 0 mV).

This type of analysis is very different than a spectral analysis, i.e. a Fourier Transform, which arbitrarily assigns combinations of sine waves of different frequencies and amplitudes, to represent the area under a curve. The assignment of the different sine waves is arbitrary and, although it gives a unique signature to a signal, there is no way to relate it to biological information, except through an empirical approach to the testing of drugs or other compounds as well as toxic chemicals.

Claims 51 - 67 and 70 - 73 are rejected under 35 USC 103(a) over Borkholder et al. in view of Georger, Jr. et al.

Borkholder et al. does not specifically disclose an intervening layer, as claimed. Furthermore, it does not disclose or suggest serum-free media. Finally, it does not inherently disclose a deconvolution analysis based on a data library, as claimed. Instead it mentions that a "pattern of spectral changes provides a unique *signature* for the compound. Matching the spectral change pattern of a test compound with the spectral change pattern for reference compounds provides a useful method for characterization of channel modulating agents." (Col. 4, lines 15 - 20, emphasis added). This passage describes using the observed changes as a fingerprinting method to associate certain channel-modulating compounds with the observed changes, but that is not the same kind of process as deconvoluting a signal to identify the

Docket No.: 7049405002
Customer No. 23,639

Application Serial No. 09/575,377

contributions of different ion channels to the signal and then to assign the test compound to one or more functional categories.

Georger Jr. et al. teaches a method of making biosensors using photolithography placement techniques. It never explicitly states that a high-impedance seal is created or that it would enable deconvolution. For instance, col. 7, lines 58 - 63 does not relate to deconvolution, but at most implies a high-impedance seal. Column 10, lines 58 - 62 describes manual placement, whereas we deposit the cells by cell culture. At col. 13, lines 19 - 24, the cell culture patterning method is to use fewer steps. The discussions at col. 16, line 45 and at col. 10, lines 50 - 65 relate to positioning.

Finally, Georger Jr. et al. uses serum (col. 17, lines 60 - 65) and states at col. 18, lines 9-10 that "adhesion is unaffected by serum." Especially where high-throughput screening is involved, the absence of serum is vital.

Claims 51- 73 are rejected under 35 USC 12, first paragraph, for written description problems. Support for the high-impedance intervening layer allowing deconvolution as claimed is found on page 11, lines 6 - 12.

Claim 51 is rejected under 35 USC 12, first paragraph, on enablement grounds. A positive deconvolution step is now recited.

Claims 51 - 73 are rejected under 35 USC 12, second paragraph, as being indefinite for failure to positively recite a deconvolution step. This has been address by amendment.

Claim 57 has been clarified with respect to the position of the insulator "surrounding" the electrode.

The dependency of claim 61 has been corrected.

Claim 67 does add a limitation over claim 66 because "known function" and "unknown function" are not the only two possibilities for DNA; some DNA has no function at all. Applicants submit the case is now in condition for allowance.

Respectfully submitted,



Robert W. Hahl, Ph.D.
Registration No. 33,893

Docket No.: 7049405002
Customer No. 23,639

Application Serial No. 09/575,377


Enclosures

Date: April 28, 2006

Patent Administrator
Bingham McCutchen LLP
Three Embarcadero Center
San Francisco, California 94111-4067
Local Telephone: (202) 778-6150
Local Facsimile: (202) 778-6155